



ELSEVIER

Available online at www.sciencedirect.comLINEAR ALGEBRA
AND ITS
APPLICATIONS

Linear Algebra and its Applications 371 (2003) 369–375

www.elsevier.com/locate/laa

On weakly unitarily invariant norm and the Aluthge transformation[☆]

K. Okubo

Mathematics Laboratory, Hokkaido University of Education, Sapporo College, 5-3-1-5 Ainosato Kitaku,
Sapporo 002-8502, Japan

Received 15 January 2003; accepted 11 February 2003

Submitted by R.A. Brualdi

Abstract

In this paper we show that

$$|||f(P^\lambda U P^{1-\lambda})||| \leq \max \{ |||f(T)|||, |||U^* \cdot f(T) \cdot U + f(0)(I - U^*U)||| \},$$

where $T \in \mathcal{B}(\mathcal{H})$, $||| \cdot |||$ is a semi-norm on $\mathcal{B}(\mathcal{H})$ which satisfies some conditions, $T = UP$ (polar decomposition), $0 \leq \lambda \leq 1$ and f is a polynomial. As a consequence of this fact, we will show that some semi-norms $||| \cdot |||$ including the ρ -radii ($0 < \rho \leq 2$) satisfy the inequality $|||f(P^\lambda U P^{1-\lambda})||| \leq |||f(T)|||$. We also give some related results.

© 2003 Published by Elsevier Inc.

AMS classification: 15A60; 47A12; 47A63

Keywords: λ -Aluthge transformation; Aluthge transformation; Weakly unitarily invariant norm; Operator radii; Numerical range

1. Introduction

We denote by $\mathcal{B}(\mathcal{H})$ the algebra of all bounded linear operators on a Hilbert space \mathcal{H} . Let $T \in \mathcal{B}(\mathcal{H})$ and $T = UP$ be a polar decomposition of T . Here P is positive semi-definite and U is partial isometry satisfying $U^*UP = P$. P is determined uniquely by $P = |T|$, but usually U is not unique for the case of $\ker(T) \neq \{0\}$ and $\ker(T^*) \neq \{0\}$. For $0 < \lambda < 1$, we define the λ -Aluthge transformation of T

[☆] Supported by Grant-Aid for Scientific Research 13640146.

E-mail address: okubo@sap.hokkyodai.ac.jp (K. Okubo).

by $P^\lambda U P^{1-\lambda}$. The λ -Aluthge transformation of T is determined uniquely without depending on any choice of U . In particular, for $\lambda = 1/2$, $\tilde{T} := P^{1/2} U P^{1/2}$ is called the *Aluthge transformation* of T (see [1]). Yamazaki [7] proved that the numerical range $W(\tilde{T})$ of the Aluthge transformation \tilde{T} of T satisfies the inclusion

$$\overline{W(\tilde{T})} \subset \overline{W(T)}, \quad (1)$$

if T admits a decomposition $T = UP$ for a unitary operator U , and Wu [8] generalized this result to the general case. The numerical radius $w(T)$ of T is defined by

$$w(T) = \sup\{|z| : z \in W(T)\}. \quad (2)$$

Eq. (1) shows

$$w(\tilde{T}) \leq w(T). \quad (3)$$

For $\rho > 0$, an operator $T \in \mathcal{B}(\mathcal{H})$ is called a ρ -contraction if there is a Hilbert space $\mathcal{K} \supset \mathcal{H}$ and a unitary operator V on \mathcal{K} such that

$$T^n = \rho Q V^n|_{\mathcal{H}} \quad (n = 1, 2, \dots). \quad (4)$$

Here Q is the orthogonal projection from \mathcal{K} to \mathcal{H} . A ρ -contraction is firstly considered by Nagy and Foiaş [6], and Holbrook defined ρ -radius $w_\rho(T)$ of T by

$$w_\rho(T) = \inf \left\{ r > 0 : \frac{1}{r} T \text{ is a } \rho\text{-contraction} \right\}. \quad (5)$$

It is known that for $0 < \rho \leq 2$, $w_\rho(\cdot)$ is a norm on $\mathcal{B}(\mathcal{H})$ but for $2 < \rho < \infty$ is only a quasi-norm. Each ρ -radius has weakly unitary invariance in the sense

$$w_\rho(VTV^*) = w_\rho(T) \quad (V : \text{unitary}). \quad (6)$$

Moreover ρ -radii have the properties:

$$w_1(T) = \|T\|,$$

where $\|\cdot\|$ is the Hilbert space operator norm and

$$w_2(T) = w(T)$$

and

$$w_\infty(T) := \lim_{\rho \rightarrow \infty} w_\rho(T) = r(T),$$

where $r(\cdot)$ is the spectral radius, that is, $r(T) = \max\{|\lambda| : \lambda \in \sigma(T)\}$. Here $\sigma(T)$ is the spectrum of T .

For references of ρ -radius see [4,6].

In this paper we treat the semi-norm $||| \cdot |||$ on $\mathcal{B}(\mathcal{H})$ satisfying the following two conditions:

$$\exists \gamma, \quad |||X||| \leq \gamma \|X\| \quad (X \in \mathcal{B}(\mathcal{H})) \quad (7)$$

and

$$|||S^*XS||| \leq \|S\|^2 \cdot |||X||| \quad (X, S \in \mathcal{B}(\mathcal{H})). \quad (8)$$

Moreover, sometimes we add the following conditions:

$$|||Q||| = 1 \quad \text{for any orthogonal projection } Q \neq 0 \quad (9)$$

and

$$\begin{aligned} XQ &= QXQ : \text{orthogonal projeciton} \\ \implies |||X||| &= \max \{ |||XQ|||, |||X(1-Q)||| \}. \end{aligned} \quad (10)$$

If $|||\cdot|||$ is equivalent to the operator norm, that is, there is a positive number γ' such that

$$\gamma' |||X||| \geq \|X\| \quad (X \in \mathcal{B}(\mathcal{H})),$$

or more naturally if

$$r(X) \leq |||X||| \quad (X \in \mathcal{B}(\mathcal{H}))$$

then the condition (9) is satisfied. For $0 < \rho \leq 2$, the ρ -radius $w_\rho(\cdot)$ is an example of a norm satisfies (7)–(10).

2. Consequences

In this paper we will give the generalization of inequality (3) to the λ -Aluthge transformation and some weakly unitarily invariant semi-norms. The following inequality is the main result in this paper.

Theorem 1. *Let $T \in \mathcal{B}(\mathcal{H})$, $T = UP$ be a polar decomposition and $|||\cdot|||$ be a semi-norm on $\mathcal{B}(\mathcal{H})$ which satisfies conditions (7) and (8). Then for $0 \leq \lambda \leq 1$ and for any polynomial f , we have the inequality*

$$|||f(P^\lambda U P^{1-\lambda})||| \leq \max \{ |||f(T)|||, |||U^* \cdot f(T) \cdot U + f(0)(I - U^*U)||| \}.$$

Proof. Denote $f(z) = f(0) + g(z)z$. Here $g(z)$ is also a polynomial. We define P_n ($n = 1, 2, \dots$) as the following:

$$P_n = \begin{cases} P & (P \text{ is invertible}), \\ P + \frac{1}{n} & (P \text{ is not invertible}). \end{cases}$$

Then P_n is invertible positive definite and

$$0 \leq P_n^\lambda - P^\lambda \leq \frac{1}{n^\lambda} \quad (n = 1, 2, \dots; 0 < \lambda \leq 1).$$

Therefore in the sense of uniform convergence we have

$$\begin{aligned} f(P^\lambda U P^{1-\lambda}) &= \lim_{n \rightarrow \infty} f(P_n^\lambda U P_n^{1-\lambda}), \\ f(T) &= \lim_{n \rightarrow \infty} f(U P_n). \end{aligned}$$

We consider the following operator-valued analytic function $\varphi_n(z)$ ($n = 1, 2, \dots$) on the strip $\mathcal{D} = \{z \in \mathcal{C} : -1/2 \leq \operatorname{Re}(z) \leq 1/2\}$ in the complex plane \mathcal{C} :

$$\varphi_n(z) \equiv f(0) + P_n^{(1/2)-z} \cdot g(UP_n) \cdot UP_n^{(1/2)+z}.$$

We will estimate the norm of $\varphi_n(z)$ on the boundary of \mathcal{D} . We have

$$\begin{aligned} \varphi_n\left(\frac{1}{2} + it\right) &= f(0) + P_n^{-it} \cdot g(UP_n)UP_n \cdot P_n^{it} \\ &= P_n^{-it} \cdot f(UP_n) \cdot P_n^{it} \end{aligned}$$

and

$$\begin{aligned} \varphi_n\left(-\frac{1}{2} + it\right) &= f(0) + P_n^{-it}P_n \cdot g(UP_n)U \cdot P_n^{it} \\ &= P_n^{-it}\{f(0) + P_n \cdot g(UP_n)U\}P_n^{it}. \end{aligned}$$

Since $P_n^{\pm it}$ is unitary, it follows from (8) that

$$\begin{aligned} \sup_{t \in \mathcal{R}} |||\varphi_n\left(\frac{1}{2} + it\right)||| &\leq |||f(UP_n)|||, \\ \sup_{t \in \mathcal{R}} |||\varphi_n\left(-\frac{1}{2} + it\right)||| &\leq |||f(0) + P_n \cdot g(UP_n)U|||. \end{aligned}$$

Therefore the three-line theorem (cf. [3, pp. 136–137]) implies that

$$\begin{aligned} |||f(P_n^\lambda UP_n^{1-\lambda})||| &= |||\varphi_n\left(\frac{1}{2} - \lambda\right)||| \\ &\leq \max\{|||f(UP_n)|||, |||f(0) + P_n \cdot g(UP_n)U|||\}. \end{aligned}$$

Taking a limit, we have by (7)

$$|||f(P^\lambda UP^{1-\lambda})||| \leq \max\{|||f(T)|||, |||f(0) + P \cdot g(T)U|||\}. \quad (11)$$

Since $U^*UP = P$,

$$\begin{aligned} f(0) + P \cdot g(T)U &= U^*\{f(0) + UP \cdot g(T)\}U + f(0)(I - U^*U) \\ &= U^* \cdot f(T) \cdot U + f(0)(I - U^*U). \end{aligned}$$

Hence we have

$$|||f(0) + P \cdot g(T)U||| = |||U^* \cdot f(T) \cdot U + f(0)(I - U^*U)||| \quad (12)$$

and by (11) and (12) Theorem 1 is proven. \square

Corollary 2. Let $T, U, P, |||\cdot|||, f, \lambda$ be the same as in Theorem 1 and at least one of the following conditions be satisfied:

- (i) $f(0) = 0$,

- (ii) $\dim(\ker(T)) \leq \dim(\ker(T^*))$,
 (iii) $||| \cdot |||$ satisfies conditions (9) and (10).

Then we have

$$|||f(P^\lambda U P^{1-\lambda})||| \leq |||f(T)|||.$$

Proof. In the case of (i) or (ii), we can easily show that

$$|||U^* \cdot f(T) \cdot U + f(0)(I - U^*U)||| = |||U^* \cdot f(T) \cdot U||| \leq |||f(T)|||.$$

In the case of (iii) and if U is not an isometry, using the conditions (9) and (10) with $Q = U^*U$, we derive

$$\begin{aligned} |||U^* \cdot f(T) \cdot U + f(0)(I - U^*U)||| &= \max \{ |||U^* \cdot f(T) \cdot U|||, |f(0)| \} \\ &\leq \max \{ |||f(T)|||, |f(0)| \}. \end{aligned} \quad (13)$$

On the other hand, since $PU^*U = P$, we obtain

$$f(T)U^*U = (f(0) + g(T)T)U^*U = f(0)U^*U + g(T)T$$

and hence

$$\begin{aligned} |f(0)| \cdot |||I - U^*U||| &= |||(I - U^*U) \cdot f(T) \cdot (I - U^*U)||| \\ &\leq |||f(T)|||. \end{aligned} \quad (14)$$

Therefore combining (13) and (14) we obtain

$$|||U^* \cdot f(T) \cdot U + f(0)(I - U^*U)||| \leq |||f(T)|||,$$

and so,

$$|||f(P^\lambda U P^{1-\lambda})||| \leq |||f(T)|||. \quad \square$$

Corollary 3. Let T , U , P , f , λ be the same as in Theorem 1. Then for $0 < \rho$ we have

$$w_\rho(f(P^\lambda U P^{1-\lambda})) \leq w_\rho(f(T)). \quad (15)$$

In particular,

$$\|f(P^\lambda U P^{1-\lambda})\| \leq \|f(T)\| \quad \text{and} \quad w(f(P^\lambda U P^{1-\lambda})) \leq w(f(T)). \quad (16)$$

Proof. For $0 < \rho \leq 2$, ρ -radius is a norm which satisfies conditions (7)–(10), and hence we can use Corollary 2(iii) to prove (15). In particular, if $\rho = 1$ and $\rho = 2$ then we have (16). We may assume $\rho > 2$. For $X \in \mathcal{B}(\mathcal{H})$ and $\rho > 1$ it is known (cf. [2]) that a necessary and sufficient condition of $w_\rho(X) \leq 1$ is $r(X) \leq 1$ and

$$\left\| \sum_{k=1}^{\infty} \frac{(\rho-1)^k |z|^{k-1}}{\rho^k} X^k \right\| \leq \rho-1 \quad \text{for } |z| \leq 1. \quad (17)$$

To prove (15) we need only check that if $w_\rho(f(T)) \leq 1$ then $w_\rho(f(P^\lambda U P^{1-\lambda})) \leq 1$. Let $w_\rho(f(T)) \leq 1$. Then by (16)

$$\begin{aligned} r(f(P^\lambda U P^{1-\lambda})) &= \lim_{n \rightarrow \infty} \|(f(P^\lambda U P^{1-\lambda}))^n\|^{1/n} \\ &\leq \lim_{n \rightarrow \infty} \|(f(T))^n\|^{1/n} = r(f(T)) \leq 1. \end{aligned}$$

Also since

$$\left\| \sum_{k=1}^N \frac{(\rho-1)^k |z|^{k-1}}{\rho^k} (f(P^\lambda U P^{1-\lambda}))^k \right\| \leq \left\| \sum_{k=1}^N \frac{(\rho-1)^k |z|^{k-1}}{\rho^k} (f(T))^k \right\|$$

for $|z| \leq 1$ and for any integer N by (16), taking a limit, we have

$$\begin{aligned} \left\| \sum_{k=1}^{\infty} \frac{(\rho-1)^k |z|^{k-1}}{\rho^k} (f(P^\lambda U P^{1-\lambda}))^k \right\| &\leq \left\| \sum_{k=1}^{\infty} \frac{(\rho-1)^k |z|^{k-1}}{\rho^k} (f(T))^k \right\| \\ &\leq \rho - 1 \quad \text{for } |z| \leq 1. \end{aligned}$$

So we can prove $w_\rho(f(P^\lambda U P^{1-\lambda})) \leq 1$. \square

Corollary 4 (cf. [5, Corollary 4.3]). *Let T, U, P, f, λ be the same as in Theorem 1. Then we have the following inclusion on the numerical range $W(\cdot)$:*

$$\overline{W(f(P^\lambda U P^{1-\lambda}))} \subset \overline{W(f(T))}.$$

Proof. In general, it is known that for $X \in \mathcal{B}(\mathcal{H})$ and each $1 \leq \rho \leq 2$,

$$\overline{W(X)} = \bigcap_{\mu \in \mathcal{C}} \{z; |z - \mu| \leq w_\rho(X - \mu)\}. \quad (18)$$

By Corollary 3, we can see that

$$w_\rho(f(P^\lambda U P^{1-\lambda}) - \mu) \leq w_\rho(f(T) - \mu).$$

Hence using (18) we have the conclusion. \square

Acknowledgements

The author would like to express his hearty thanks to T. Ando, M. Ito, H. Nakazato and T. Yamazaki for helpful discussions.

References

- [1] A. Aluthge, On p -hyponormal operators for $0 < p < 1$, *Integral Equations Operator Theory* 13 (1990) 307–315.
- [2] T. Ando, K. Okubo, Hölder type inequalities associated with operator radii and Schur products, *Linear and Multilinear Algebra* 43 (1997) 53–61.

- [3] I.C. Gohberg, M.G. Krčín, Introduction to the theory of linear nonselfadjoint operators, Amer. Math. Soc. 18 (1969) Translations of Mathematical Monographs.
- [4] J.A.R. Holbrook, On the power-bounded operators of Sz.-Nagy and Foiaş, Acta Sci. Math. (Szeged) 29 (1968) 297–310.
- [5] M. Ito, H. Nakazato, K. Okubo, T. Yamazaki, On generalized numerical range of the Aluthge transformation, Linear Algebra Appl., in press.
- [6] Sz. Nagy, C. Foiaş, Harmonic analysis of operators on Hilbert space, North-Holland, Amsterdam, 1970.
- [7] T. Yamazaki, On numerical range of the Aluthge transformation, Linear Algebra Appl. 341 (2002) 111–117.
- [8] P.Y. Wu, Numerical range of Aluthge transform of operator, Linear Algebra Appl. 357 (2002) 295–298.